

Coherence of Supernova Neutrinos

Jörn Kersten


University of Hamburg



Based on work done in collaboration with A.Yu. Smirnov

Interesting Effects with Supernova Neutrinos

- Oscillations inside the Earth
- MSW and **collective effects** inside the supernova




Is **coherence** preserved in these cases?

Decoherence

- Coherent 2-flavor oscillation probability

$$P(\nu_e \rightarrow \nu_e) = |\cos^2 \theta + \sin^2 \theta e^{i\phi}|^2 = 1 - \sin^2 2\theta \sin^2 \frac{\Delta m^2 L}{4E}$$

- Oscillation phase $\phi = -\frac{\Delta m^2 L}{2E}$
- Mass eigenstates have different velocities
 $\leadsto$ Wave packets cease to overlap



- $\leadsto$ Probability is **incoherent** sum

$$P(\nu_e \rightarrow \nu_e) = |\cos^2 \theta|^2 + |\sin^2 \theta e^{i\phi}|^2 = \cos^4 \theta + \sin^4 \theta$$

- $\leadsto$ **No oscillations** between supernova and Earth

Coherence Length of Supernova Neutrinos

- Wave packets overlap up to **coherence length** $L_{\text{coh}} \sim \frac{\sigma}{\Delta v}$
- Depends on
 - Size of wave packets σ
 - Velocity difference Δv of mass eigenstates

Size of Wave Packets

- Determined by length of production process
- Electron capture $pe^- \rightarrow n\nu_e$
- Time scale $\sim$ time electron needs to cross proton, $\tau \sim \sigma_e/v_e$
- Temperature $\sim 5 \text{ MeV} \Rightarrow$ electron relativistic, $v_e \sim 1$
- Size of electron wave packet $\sigma_e \sim$ mean free path

Size of Wave Packets

- Determined by length of production process
- Electron capture $pe^- \rightarrow n\nu_e$
- Time scale $\sim$ time electron needs to cross proton, $\tau \sim \sigma_e/v_e$
- Temperature ~ 5 MeV $\Rightarrow$ electron relativistic, $v_e \sim 1$
- Size of electron wave packet $\sigma_e \sim$ mean free path
- Result: $\sigma \sim \sigma_e \sim (4\pi\alpha^2 n)^{-1/3} \sim 10^{-11}$ cm
- For comparison:
 - Atmospheric neutrinos: $\sigma \sim 1$ cm
 - Reactor neutrinos: $\sigma \sim 10^{-8}$ cm
- $\sigma_E \sim 1/\sigma \sim 1$ MeV not much smaller than $E \sim 10$ MeV

Coherence Length of Supernova Neutrinos

- Wave packets overlap up to **coherence length** $L_{\text{coh}} \sim \frac{\sigma}{\Delta v}$
- Depends on
 - Size of wave packets $\sigma \sim 10^{-11}$ cm
 - Velocity difference Δv of mass eigenstates

Coherence Length of Supernova Neutrinos

- Wave packets overlap up to **coherence length** $L_{\text{coh}} \sim \frac{\sigma}{\Delta v}$
- Depends on
 - Size of wave packets $\sigma \sim 10^{-11}$ cm
 - Mass difference $\Delta m_{\text{atm}}^2 \sim 10^{-3} \text{ eV}^2$ or $\Delta m_{\text{sol}}^2 \sim 10^{-5} \text{ eV}^2$
 - Neutrino energy $E \sim 10 \text{ MeV}$

Coherence Length of Supernova Neutrinos

- Wave packets overlap up to **coherence length** $L_{\text{coh}} \sim \frac{\sigma}{\Delta v}$
- Depends on
 - Size of wave packets $\sigma \sim 10^{-11}$ cm
 - Mass difference $\Delta m_{\text{atm}}^2 \sim 10^{-3} \text{ eV}^2$ or $\Delta m_{\text{sol}}^2 \sim 10^{-5} \text{ eV}^2$
 - Neutrino energy $E \sim 10 \text{ MeV}$
 - Matter density
 - Mixing angle $\theta_{13} \sim 9^\circ$ or $\theta_{12} \sim 30^\circ$

Coherence Length of Supernova Neutrinos

- Wave packets overlap up to **coherence length** $L_{\text{coh}} \sim \frac{\sigma}{\Delta v}$
- Depends on
 - Size of wave packets $\sigma \sim 10^{-11}$ cm
 - Mass difference $\Delta m_{\text{atm}}^2 \sim 10^{-3} \text{ eV}^2$ or $\Delta m_{\text{sol}}^2 \sim 10^{-5} \text{ eV}^2$
 - Neutrino energy $E \sim 10 \text{ MeV}$
 - Matter density
 - Mixing angle $\theta_{13} \sim 9^\circ$ or $\theta_{12} \sim 30^\circ$
- Supernova, inner region:
 $L_{\text{coh}} \sim 100 \text{ km} \lesssim \text{length scale of collective effects}$
- Earth: $L_{\text{coh}} \sim 1000 \text{ km}$

Coherence Length of Supernova Neutrinos

- Wave packets overlap up to **coherence length** $L_{\text{coh}} \sim \frac{\sigma}{\Delta v}$
- Depends on
 - Size of wave packets $\sigma \sim 10^{-11}$ cm
 - Mass difference $\Delta m_{\text{atm}}^2 \sim 10^{-3} \text{ eV}^2$ or $\Delta m_{\text{sol}}^2 \sim 10^{-5} \text{ eV}^2$
 - Neutrino energy $E \sim 10 \text{ MeV}$
 - Matter density
 - Mixing angle $\theta_{13} \sim 9^\circ$ or $\theta_{12} \sim 30^\circ$
- Supernova, inner region:
 $L_{\text{coh}} \sim 100 \text{ km} \lesssim$ **length scale of collective effects**
- Earth: $L_{\text{coh}} \sim 1000 \text{ km}$

Supernova neutrinos are special

Very short wave packets $\leadsto$ short coherence length

Sometimes Decoherence Is Irrelevant

$$\begin{aligned} P(\nu_e \rightarrow \nu_e) &= |\cos^2 \theta + \sin^2 \theta e^{i\phi}|^2 \\ &= \cos^4 \theta + \sin^4 \theta + 2 \cos^2 \theta \sin^2 \theta \cos \phi \end{aligned}$$

- Measured probability: average over **energy resolution** ΔE of detector

Sometimes Decoherence Is Irrelevant


$$\begin{aligned} P(\nu_e \rightarrow \nu_e) &= |\cos^2 \theta + \sin^2 \theta e^{i\phi}|^2 \\ &= \cos^4 \theta + \sin^4 \theta + 2 \cancel{\cos^2 \theta \sin^2 \theta \cos \phi} \end{aligned}$$

- Measured probability: average over **energy resolution** ΔE of detector
- ΔE too large $\leadsto$ interference term vanishes

Sometimes Decoherence Is Irrelevant

$$\begin{aligned}P(\nu_e \rightarrow \nu_e) &= |\cos^2 \theta + \sin^2 \theta e^{i\phi}|^2 \\&= \cos^4 \theta + \sin^4 \theta + 2 \cos^2 \theta \sin^2 \theta \cos \phi\end{aligned}$$

- Measured probability: average over **energy resolution** ΔE of detector
- ΔE too large $\leadsto$ interference term vanishes
 $\leadsto$ Need sufficient energy resolution to observe oscillations



Sometimes Decoherence Is Irrelevant

$$\begin{aligned} P(\nu_e \rightarrow \nu_e) &= |\cos^2 \theta + \sin^2 \theta e^{i\phi}|^2 \\ &= \cos^4 \theta + \sin^4 \theta + 2 \cos^2 \theta \sin^2 \theta \cos \phi \end{aligned}$$

- Measured probability: average over **energy resolution** ΔE of detector
- ΔE too large $\leadsto$ interference term vanishes
 $\leadsto$ Need sufficient energy resolution to observe oscillations
- Corresponding uncertainty of arrival **time** $\Delta t \sim 1/\Delta E$
- Coherence preserved for wave packets arriving within Δt , even if they are spacially separated
 $\leadsto$ **Detector restores coherence**


Sometimes Decoherence Is Irrelevant

$$\begin{aligned} P(\nu_e \rightarrow \nu_e) &= |\cos^2 \theta + \sin^2 \theta e^{i\phi}|^2 \\ &= \cos^4 \theta + \sin^4 \theta + 2 \cos^2 \theta \sin^2 \theta \cos \phi \end{aligned}$$

- Measured probability: average over **energy resolution** ΔE of detector
- ΔE too large $\leadsto$ interference term vanishes
 $\leadsto$ Need sufficient energy resolution to observe oscillations
- Corresponding uncertainty of arrival **time** $\Delta t \sim 1/\Delta E$
- Coherence preserved for wave packets arriving within Δt , even if they are spacially separated
 $\leadsto$ **Detector restores coherence**
- Always the case in vacuum and matter with slowly changing density (adiabatic case)
- Does this change in **non-adiabatic** case?


Adiabaticity Violation

- Simplest case: density step
- Each wave packet **splits up** into two




Adiabaticity Violation

- Simplest case: density step
- Each wave packet **splits up** into two




Adiabaticity Violation

- Simplest case: density step
- Each wave packet **splits up** into two




Adiabaticity Violation

- Simplest case: density step
- Each wave packet **splits up** into two



Adiabaticity Violation


- Simplest case: density step
- Each wave packet **splits up** into two



Adiabaticity Violation

- Simplest case: density step
- Each wave packet **splits up** into two
- Different oscillation phase in each layer
- Complete coherence:

$$P = \left| a + b e^{i\phi_1} + c e^{i\phi_2} + \dots \right|^2$$



Adiabaticity Violation

- Simplest case: density step
- Each wave packet **splits up** into two
- Different oscillation phase in each layer
- Complete coherence:

$$P = \left| a + b e^{i\phi_1} + c e^{i\phi_2} + \dots \right|^2$$


- Complete decoherence:

$$P = a^2 + b^2 + c^2 + \dots$$

- Energy resolution good enough to resolve **all** oscillation features
⇒ detector restores coherence as before

Incomplete Averaging

$$P = \left| a + b e^{i\phi_1} + c e^{i\phi_2} + \dots \right|^2$$
$$= a^2 + b^2 + c^2 + \dots + 2ab \cos \phi_1 + 2bc \cos(\phi_2 - \phi_1)$$




Energy resolution may be

- **too bad** to observe $\cos \phi_1$ term
- **good enough** to observe $\cos(\phi_2 - \phi_1)$ term if $\phi_1 \approx \phi_2$

Incomplete Averaging

$$\begin{aligned} P &= \left| a + b e^{i\phi_1} + c e^{i\phi_2} + \dots \right|^2 \\ &= a^2 + b^2 + c^2 + \dots + 2ab \cos \phi_1 + 2bc \cos(\phi_2 - \phi_1) \end{aligned}$$




Energy resolution may be

- **too bad** to observe $\cos \phi_1$ term
- **good enough** to observe $\cos(\phi_2 - \phi_1)$ term if $\phi_1 \approx \phi_2$

But only if wave packets overlap (coherent case)?


Wave Packet Catch-Up

- Wave packets don't need to separate forever:




Wave Packet Catch-Up

- Wave packets don't need to separate forever:




Wave Packet Catch-Up

- Wave packets don't need to separate forever:



Wave Packet Catch-Up

- Wave packets don't need to separate forever:



Wave Packet Catch-Up

- Wave packets don't need to separate forever:
- Wave packets catch up $\iff \Delta E$ good enough to observe $\cos(\phi_2 - \phi_1)$ term
- Decoherence and energy averaging are **equivalent**

Wave Packet Catch-Up

- Wave packets don't need to separate forever:
- Wave packets catch up $\longleftrightarrow$ ΔE good enough to observe $\cos(\phi_2 - \phi_1)$ term
- Decoherence and energy averaging are **equivalent**
- Open questions:
 - How general is this?
 - Are collective effects in a supernova different?

Effect of Wave Packet Size on Spectrum

- Finite $\sigma_E \leadsto$ symmetric broadening of energy spectrum $f(E)$
- Change relevant for $\sigma_E^2 \gtrsim f/f''$

Effect of Wave Packet Size on Spectrum

- Finite $\sigma_E \leadsto$ symmetric broadening of energy spectrum $f(E)$
- Change relevant for $\sigma_E^2 \gtrsim f/f''$
- Example:
 - $f(E) \propto E^\alpha \exp[-(\alpha + 1) E/\bar{E}]$
 - Mean energy $\bar{E} = 10 \text{ MeV}$
 - $\alpha = 3$
 - $\sigma_E = 2 \text{ MeV}$
- $\leadsto$ **Potentially observable** deformation of spectrum for $E \lesssim 3 \text{ MeV}$ and around $E \simeq 5 \text{ MeV}$

Supernova Neutrinos

- Extremely **small** size of **wave packets** $\sigma \sim 10^{-11}$ cm
- No experimental consequences in simple examples
- Possibly relevant for
 - **collective effects** in a supernova
 - energy **spectrum**