THEORY #2

Statistical Method & Techniques for Data Analysis

Erasmus+ EU programme	Alexis Pompili (*) University of Bari Aldo Moro	
Erasmus ⁺ Teaching Mobility Program / 2025 May 6-13 @ NKUA (Physics Dpt.)		



PART 2A - CORE

A.Pompili (E+)

Statistical M&T for Data Analysis

PROBABILITY DENSITY FUNCTIONS

Probability distribution function (aka p.d.f.): distribution of the probability for a RV to assume a certain value among those allowed

In other words: the p.d.f. of a RV is the law which rules the assumption of a certain value by the RV in one measurement/experiment

We will see during this course that: the link between experiment and theoretical model indeed happens through the p.d.f., that is predicted by the model to describe (the result of) an experiment

Consider a discrete random variable x having more than one possible elementary result, that is (x₁, ..., x_N) each occurring with a probability P(x_i), where i = 1, ..., N, thus associated to each of the possible results.
The function that associates the probability P(x_i) to each possible value x_i is called probability distribution.
Note : the result of an event is not predictable but - instead - the probability distribution of the results can be known.

Probability distribution function (aka p.d.f.): distribution of the probability for a RV to assume a certain value among those allowed

In other words: the p.d.f. of a RV is the law which rules the assumption of a certain value by the RV in one measurement/experiment

We will see during this course that: the link between experiment and theoretical model indeed happens through the p.d.f., that is predicted by the model to describe (the result of) an experiment

Consider a discrete random variable x having more than one possible elementary result, that is (x₁, ..., x_N) each occurring with a probability P(x_i), where i = 1, ..., N, thus associated to each of the possible results.
The function that associates the probability P(x_i) to each possible value x_i is called probability distribution.
Note : the result of an event is not predictable but - instead - the probability distribution of the results can be known.

The probability of a random event E corresponding to a set of distinct possible elementary results $(x_{E_1}, ..., x_{E_K})$ where $x_{E_j} \in \Omega = (x_1, ..., x_N)$ for all j = 1, ..., K, is, according to the 3rd Kolmogorov's axiom, given by:

$$P\left(\bigcup_{j=1}^{K} \{x_{E_j}\}\right) = P(\{x_{E_1}, \dots, x_{E_K}\}) = P(E) = \sum_{j=1}^{K} P(x_{E_j})$$

From the 2nd Kolmogorov's axiom, the probability of the event Ω corresponding to the set of all possible values must be: $\sum_{i=1}^{N} P(x_i) = 1$

From the 1st Kolmogorov's axiom:
$$P(x_{E_j}) \ge 0 \quad \forall j \Longrightarrow P(E \subset \Omega) \ge 0$$

i=1 (normalization condition)

A.Pompili (E+)

Statistical M&T for Data Analysis.

Probability Density Function (p.d.f.) - II

 $\mathbf{\Sigma}$ Most quantities of interest to us are continuous, thus we will treat mainly the continuous case. The discrete probability introduced in the previous slide can be generalized to the continuous case with the replacement ... $\sum_{n=1}^{\infty} \Rightarrow \int_{\Omega} dn$

In the discrete case we deal with a genuine probability function; in the continuous case we must introduce a probability density function!

Let us consider a sample space $\Omega \subseteq \mathbb{R}^n$. Each random experiment will lead to a measurement corresponding to one point $\vec{x} \in \Omega$. $\mathbf{\Sigma}$ We can associate a probability density $f(\vec{x}) = f(x_1, ..., x_n)$ to any point $\vec{x} \in \Omega$. Of course, $f(\vec{x}) \ge 0$ (1st axiom).

The probability of an event A with $A \subseteq \Omega$, namely the probability that $\vec{x} \in A$ is given by $P(A) = \int_A f(x_1, ..., x_n) d^n x$

The function $f(\vec{x})$ is called **probability density function p.d.f.** ! The function $f(x_1, ..., x_n)d^nx$ can be interpreted as differential probability.

The normalization condition can be expressed as: $\int_{\Omega} f(x_1, \dots, x_n) d^n x = 1$

 $\mathbf{\Sigma}$

Probability Density Function (p.d.f.) - II

Most quantities of interest to us are continuous, thus we will treat mainly the continuous case. $\mathbf{\Sigma}$ The discrete probability introduced in the previous slide can be generalized to the continuous case with the replacement ... $\sum_{n=1}^{\infty} \Rightarrow \int_{\Omega}$

In the discrete case we deal with a genuine probability function; in the continuous case we must introduce a probability density function!

Let us consider a sample space $\Omega \subseteq \mathbb{R}^n$. Each random experiment will lead to a measurement corresponding to one point $\vec{x} \in \Omega$. $\mathbf{\Sigma}$ We can associate a probability density $f(\vec{x}) = f(x_1, ..., x_n)$ to any point $\vec{x} \in \Omega$. Of course, $f(\vec{x}) \ge 0$ (1st axiom).

The probability of an event A with $A \subseteq \Omega$, namely the probability that $\vec{x} \in A$ is given by $P(A) = \int_A f(x_1, ..., x_n) d^n x$

The function $f(\vec{x})$ is called **probability density function p.d.f.** ! The function $f(x_1, ..., x_n)d^nx$ can be interpreted as differential probability.

The normalization condition can be expressed as: $\int_{\Omega} f(x_1, \dots, x_n) d^n x = 1$

In 1 dim: Probability of the outcome X to be within the continuous interval of possible values [x, x + dx] is $P(x \le X \le x + dx) = f(x) \cdot dx$ The **p.d.f.** f(x) is of course normalized by the condition : $\int_{-\infty}^{+\infty} f(x) dx = 1$

It can be verified that :

the p.d.f. corresponds to an histogram of the RV x normalized to the unity area in the limit for which ... - the bin width $\rightarrow 0$ - the total # of entries $\rightarrow \infty$

Cumulative Distribution Function (c.d.f.)

The cumulative distribution function (c.d.f.) is the probability that the value of a r.v. will be \leq a specific value. The c.d.f. is denoted by the capital letter corresponding to the small letter signifying the p.d.f. The c.d.f. is thus given by

$$F(x) = \int_{-\infty}^{x} f(x') \, \mathrm{d}x' = P(X \le x)$$

Clearly, $F(-\infty) = 0$ and $F(+\infty) = 1$.

Properties of the c.d.f.:

- $0 \le F(x) \le 1$
- F(x) is monotone and not decreasing.
- $P(a \le X \le b) = F(b) F(a)$
- F(x) discontinuous at x implies

$$P(X = x) = \lim_{\delta x \to 0} \left[F(x + \delta x) - F(x - \delta x) \right]$$
, *i.e.*, the size of the jump.

F(x) continuous at x implies P(X = x) = 0.

The c.d.f. can be considered to be more fundamental than the p.d.f. since the c.d.f. is an actual probability rather than a probability density. However, in applications we usually need the p.d.f. Sometimes it is easier to derive first the c.d.f. from which you get the p.d.f. by

$$f(x) = \frac{\partial F(x)}{\partial x} \tag{2.4}$$

Note: the p.d.f. for *F* is **uniformly distributed** in [0,1]: $\frac{dP}{dF} = \frac{dP}{dx} \cdot \frac{dx}{dF} = \frac{f(x)}{f(x)} = 1$



A.Pompili (E+)

Statistical M&T for Data Analysis.

Library of p.d.f.s in ROOT/RooFit

RooFit provides a collection of compiled standard PDF classes



Easy to extend the library: each p.d.f. is a separate C++ class

Attributes of a p.d.f. : mode & median



Expectation value of a p.d.f. (sometimes called "*Mean*" which is very misleading actually! Better *population mean*): represents the central value of a p.d.f. and it is defined as: $\mu = E[x] = \int_{-\infty}^{+\infty} x f(x) dx$

Note: E[x] is not a function of x (there is an integral on x !) but depends on the distribution of the values taken by x (that is on the shape of the p.d.f.)

The mean is often a good measure of location, *i.e.*, it frequently tells roughly where the most probable region is, but not always.



Properties: $a = \cot \Rightarrow E[a] = a \& E[ax] = a \cdot E[x]$ if **u** is a function of **x**: $E[au(x)] = a \cdot E[u(x)]$ where $E[u(x)] = \int_{-\infty}^{+\infty} u(x) f(x) dx$

E is a linear operator: $E[a_1u(x) + a_2v(x)] = a_1E[u(x)] + a_2E[v(x)]$

Attributes of a **p.d.f.** : example of the Maxwell-Boltzmann distributuion



Example: distribution of the squared velocity of the gas molecules exiting the hole of a cavity/container

For this distribution:

the expectation value ("Mean") > Median



(note: this is the effect of the large tail on the right)

⁻⁻⁻⁻⁻ Median > Mode

Attribute of a c.d.f. : quantile of order α

A useful concept related to the cumulative distribution is the so-called **quan**tile of order α or α -point. The quantile x_{α} is defined as the value of the random variable x such that $F(x_{\alpha}) = \alpha$, with $0 \le \alpha \le 1$. That is, <u>the quantile is simply</u> the inverse function of the cumulative distribution,

$$x_{\alpha} = F^{-1}(\alpha). \tag{1.17}$$

$$\int_{-\infty}^{x_{\alpha}} f(x) dx = \alpha = 1 - \int_{x_{\alpha}}^{+\infty} f(x) dx$$

A commonly used special case is $x_{1/2}$, called the **median** of x. This is often used as a measure of the typical 'location' of the random variable, in the sense that there are equal probabilities for x to be observed greater or less than $x_{1/2}$.



The moments are particular expectation values. The moments of order **m** are defined as: $E[x^m] = \int_{-\infty}^{+\infty} x^m f(x) dx$. Therefore: **moment of order 1 = expectation value**

> It is possible to introduce also the <u>central</u> moments of order m, defined as: $E[(x - \mu)^m] = \int_{-\infty}^{+\infty} (x - \mu)^m f(x) dx$.

Note: if μ is finite ... the central moment of order 1 is null for any μ : =1 (normalization) $E[(x - \mu)^{m=1}] = \int_{-\infty}^{+\infty} (x - \mu)f(x)dx = \int_{-\infty}^{+\infty} xf(x)dx - \mu \int_{-\infty}^{+\infty} f(x)dx = \int_{-\infty}^{+\infty} xf(x)dx - \mu = E[x] - \mu = \mu - \mu = 0$

Note also: if f(x) is symmetric ... the central moments of odd orders (m = 1, 3, 5, ...) are null !

The central moment of order 2 is called variance and represents the spread of the f(x) around the expectation value. See details next slide!

Attribute of a p.d.f. : variance

> Variance of a p.d.f. is defined as:

$$\sigma_x^2 = V[x] = E[(x - \mu)^2] = \int_{-\infty}^{+\infty} (x - \mu)^2 f(x) dx = \mu = 1 \text{ (norm.)}$$
$$= \int_{-\infty}^{+\infty} x^2 f(x) dx - 2\mu \int_{-\infty}^{+\infty} x f(x) dx + \mu^2 \int_{-\infty}^{+\infty} f(x) dx dx = E[x^2] - 2\mu^2 + \mu^2 = E[x^2] - \mu^2 = E[x^2] - (E[x])^2$$

The squared root of the variance is called standard deviation of x and denoted by σ_x . It is often useful because it has the same dimentional units of x and thus ...

... it represents the *spread* of the p.d.f. around its expectation value.

Property: $V[ax] = a^2 \cdot V[x]$, with a = cost.

Indeed: $V[ax] = E[a^2x^2] - (E[ax])^2 = a^2 E[x^2] - (aE[x])^2 = a^2 \cdot (E[x^2] - (E[x])^2) = a^2 \cdot V[x]$

Note: other attibutes like skewness (asymmetry indicator) and kurtosis (sharpness indicator) are defined in the in-depth part.

Often a data sample under analysis is the sum of two (or more) subsamples distributed according to different p.d.f.s; $\mathbf{\Sigma}$ an obvious example is the sum of a certain signal and one (or more) background(s).

Let's express the fractions of events belonging to each subsample as φ_i and the $f_i(x)$ are the corresponding p.d.f.s of the r.v. x; then:

overall p.d.f. : $f(x) = \sum_{i} \varphi_{i} f_{i}(x)$

An obvious example is ($i \equiv 1$ for signal, $i \equiv 2$ for background): $f_{tot}(x) = \varphi_{sig} f_{sig}(x) + \varphi_{bkg} f_{bkg}(x) = \varphi_{sig} f_{sig}(x) + (1 - \varphi_{sig}) f_{bkg}(x)$

The (overall) expectation value of x is (where μ_i represents the expectation value of x for each subsample):

 $\mu \equiv E[x] = \int_{-\infty}^{+\infty} x f(x) dx = \sum_{i} \varphi_{i} \cdot \int_{-\infty}^{+\infty} x f_{i}(x) dx = \sum_{i} \varphi_{i} E_{i}[x] \equiv \sum_{i} \varphi_{i} \mu_{i} \quad \text{of the expectation values weighted by}$

the overall expectation value is the mean the relative fractions in the mixture

For the variance see next slide!

 \rightarrow The variance of a r.v. x for a mixture of subsamples is:

$$V[x] = E[(x - \mu)^2] = \sum_i \varphi_i \cdot E_i [(x - \mu)^2] \quad \text{where} \quad \mu = \sum_i \varphi_i \mu_i \quad \text{is the expectation value over the mixture}$$

To get a more familiar expression we must introduce the deviations $\delta_i = (\mu - \mu_i)$ and introduce in the upper expression $\mu = \delta_i + \mu_i$;

with some algebra (reported in a slide in the in-depth part) we can get:

$$V[x] = \sum_{i} \varphi_{i} \cdot \left\{ V_{i}[x] + \left[\sum_{j \neq i} \varphi_{i}(\mu_{j} - \mu_{i}) \right]^{2} \right\} \ge \mathbf{0}$$

generally, the variance is **not** just the simple mean of the variances of the sub-samples weighted by the relative fractions in the mixture, since it is always augmented because of the fact that sub-samples can have different expectation values

>> On the other hand, ...

(\$)
$$V[x] = \sum_{i} \varphi_{i} \cdot V_{i}[x]$$
 IF $\mu_{i} = \mu \forall i$

IF all the distributions of r.v. x for each sub-sample in the mixture are characterized by the same expectation value ...
 ... the overall variance is the mean of the variances weighted by relative fractions in the mixture

(see an example in the following slide)

 Σ Example: suppose to reconstruct with the CMS detector the dimuon decays of the charmonium state $\psi(2S)$: $\psi(2S) \rightarrow \mu^+ \mu^-$



Suppose to put together two signal subsamples of $\psi(2S)$ candidates, one with $y \in [0., 0.2]$ and the other with $y \in [1.4, 1.6]$ [we neglect the combinatorial background of 2 μ s (pairs by random combinations)]. The r.v. represented by the reconstructed mass, $m(\mu\mu)$, is characterized, in the two sub-samples, by the same expectation **value** [the mass of the $\psi(2S)$]; instead, the two standard deviations (square root of the two variances), that represent the mass resolution, are different for the two sub-samples since the mass resolution depends on the quality of the track reconstruction of the two μ s which - in turn - depend on the detection technology of the μ -chambers: the DTs ensures a better quality w.r.t. the CSCs.

Statistical M&T for Data Analysis.

Mixture of subsamples - example - II



FUNCTIONS of a R.V.

A.Pompili (E+)

Statistical M&T for Data Analysis

Function of a random variable - I

Often experimentalists carry out indirect measurements, i.e the observable of interest is a function of direct measurements. For this reason we need to introduce functions of random variables! First of all have in mind that: functions of random variables are random variables themselves !

Suppose u(x) is a continuous function of a continuous random variable x distributed according to the p.d.f. f(x).

The question now is: what is the p.d.f. $\mathbf{g}(\mathbf{u})$ that describes the distribution of $\mathbf{u}(\mathbf{x})$?

It is possible to answer requiring that the probability of x to assume values between x and x + dx has to be equal to the probability for u(x) to get values between u and u + du. If the fuction u(x) can be inverted to obtain x(u) and the trasformation is 1-to-1 (i.e. bijective)... ... then we can write:

u(x)

4

2

du

dx

>> We can develop in series the u(x) in an interval of x around μ ; thus we can substitute u(x) with its development in series and for simplicity we can stop to the 2nd order:

$$u(x) \rightarrow \left. \frac{\partial u}{\partial x} \right|_{x \sim \mu} \cdot (x - \mu) + \left. \frac{1}{2!} \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \left|_{x \sim \mu} + \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \left|_{x \sim \mu} + \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \left|_{x \sim \mu} + \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \left|_{x \sim \mu} + \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \left|_{x \sim \mu} + \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \left|_{x \sim \mu} + \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \left|_{x \sim \mu} + \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \left|_{x \sim \mu} + \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \left|_{x \sim \mu} + \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \left|_{x \sim \mu} + \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \left|_{x \sim \mu} + \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \left|_{x \sim \mu} + \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \left|_{x \sim \mu} + \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \left|_{x \sim \mu} + \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot (x - \mu)^2 + \sum_{x \sim \mu} \frac{\partial^2 u}{\partial x^2} + \sum$$

The substitution is applied inside the expression $E[u(x)] = \int_{-\infty}^{+\infty} u(x) f(x) dx$...and after a bit of algebra one gets:

$$E[u(x)] \cong \left[E[u(\mu)] + \frac{1}{2} \frac{\partial^2 u}{\partial x^2} \right|_{x \sim \mu} \cdot V[x]$$
$$= u(\mu)$$

<u>Conclusions</u>: 1) unless V[x] = 0 ... the expectation value of u(x) is not equal to the value of the function calculated with the expectation value of x, namely $u(\mu) \rightarrow E[u]$

 $\implies E[u(x)] \neq u(\mu)$

2) if u(x) is a **linear function** of x ther $E[u(x)] = u(\mu)$

3) if $\frac{\partial^2 u}{\partial x^2}\Big|_{x \sim \mu}$ is small (slowly-varying shape) this equality holds with a good approximation: $E[u(x)] \approx u(\mu)$

Dealing with more than one R.V. : MARGINAL & CONDITIONAL PDFs

Case of more than 1 random variable

> If the measurement is characterized not by just one observable but instead by more than one it means we have to deal with more than 1 random variable and specifically with a vector of random variables $\vec{x} = (x_1, ..., x_N)$; the associated p.d.f. would be $f(\vec{x})$. Its meaning is as follows:

for an infinitesimal volume centered on \vec{x} of sides $dx_1, ..., dx_N$ that we label as $I_{\vec{x}, d\vec{x}}$, the associated probability can be expressed as ...

 $P(\vec{X} \in I_{\vec{x}, d\vec{x}}) = f(\vec{x}) d\vec{x}$

We will discuss the easiest case of two r.v.s in the net slides!

> In general, this will be a complicated multi-dimentional, unless $x_1, ..., x_N$ are <u>all independent</u> among each other ... and in this particular case the expression of $f(\vec{x})$ is the following product:

$$f(\vec{x}) = \prod_{i} f_i(x_i)$$
 (where f_i is the p.d.f. of x_i)

We will come back to this possible *factorization* soon, in next slides!

Two random variables - joint p.d.f.

> Let's consider - in the following - to deal with only 2 random variables: $x \otimes y$! Let's also continue to imagine to be working in the infinite sample assumption (infinite points (x, y) in the plot): we deal with an (infinite) population, not a finite sample!

As depicted in the *scatter plot* in the figure, we consider :

Event A (vertical narrow band): observe x-values in [x, x + dx] and y-values everywhere

Event **B** (horizontal narrow band): observe y-values in [y, y + dy] and x-values everywhere

The event $A \cap B$ is associated to the intersection of the two bands. Its associated probability can be expressed in terms of a joint p.d.f. (corresponding to the **density of points**) :

$$P(A \cap B) = P(x \in [x, x + dx], y \in [y, y + dy]) = \frac{f(x, y)dx}{f(x, y)dx} \frac{dy}{dy}$$

The relative normalization condition can be expressed as: $\iint_{\Omega} f(x, y) \, dx \, dy = 1$



Two random variables & marginal p.d.f. - I

Suppose we want to know the probability for the r.v. x to get values in the interval [x, x + dx] independently from the value taken by the other r.v. y, i.e. we want to know the **probability of event A** (the vertical band in the scatter plot).

The band can be considered as the set of N squares of area $dxdy_i$ with the running index exhausting the full band:

$$P(A) = \sum_{i} f(x, y_{i}) dy_{i} dx \equiv f_{x}(x) dx$$



In the limit of infinitesimal all equal intervals one gets $dy_i = dy$ and the sum becomes an integral ($\sum_i dy_i \to \int_{-\infty}^{+\infty} dy$)

We can now introduce the concept of ... marginal p.d.f. which is the p.d.f. of 1 only random variable once the dependency from the other(s) is eliminated via integration of the joint p.d.f.

marginal p.d.f. in
$$x$$
: $f_x(x) = \int_{-\infty}^{+\infty} f(x, y) dy$, marginal p.d.f. in y : $f_y(y) = \int_{-\infty}^{+\infty} f(x, y) dx$

Note: the 2 marginal p.d.f.s correspond to the *normalized* functions obtained by *projection* of the scatter plot on the *x*, *y* axes (again - implicitely - in the limit of infinite entries in the scatter plot) [see next slide].

Two random variables & marginal p.d.f. - II



Statistical M&T for Data Analysis.

It is now possible to introduce the concept of **conditional p.d.f.** exploiting the definition of conditional probability :

Probability for r.v.
$$y$$
 to get values in the interval $[y, y + dy]$
for any value taken by the r.v. x (event B),
once it happened that x has got values in the interval $[x, x + dx]$
for any value taken by the r.v. y (event A)
is given by... $P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{f(x, y)dx dy}{f_x(x)dx}$
marginal p.d.f.

At this point it makes sense to introduce the...

conditional p.d.f. associated to the r.v. y given the r.v. x

(function of the y only since x has taken a specific value) as ...

$$\mathbf{h}(\mathbf{y}|\mathbf{x}) = \frac{\mathbf{f}(\mathbf{x}, \mathbf{y})}{\mathbf{f}_{\mathbf{x}}(\mathbf{x})} = \frac{\mathbf{f}(\mathbf{x}, \mathbf{y})}{\int_{-\infty}^{+\infty} f(\mathbf{x}, \mathbf{y}') d\mathbf{y}'}$$

In other words: the **conditional p.d.f. of** *y* is defined starting from the joint p.d.f. in which *x* has taken a specific vaue (thus, it is constant), <u>renormalized</u> so that it has unit area when integrating on *y* only) (always - implicitly - in the limit of infinite entries in the scatter plot)

Similar considerations exchanging the role of x and y brings to:

$$\mathbf{g}(\boldsymbol{x}|\boldsymbol{y}) = \frac{\boldsymbol{f}(\boldsymbol{x},\boldsymbol{y})}{\boldsymbol{f}_{\boldsymbol{y}}(\boldsymbol{y})} = \frac{\boldsymbol{f}(\boldsymbol{x},\boldsymbol{y})}{\int_{-\infty}^{+\infty} f(\boldsymbol{x}',\boldsymbol{y}) d\boldsymbol{x}'}$$

Two random variables & Conditional p.d.f. - II



The conditional p.d.f.s can be easily represented as *normalized* projections of narrow bands (large dx) in the conditioning variable

[borrowed by Cowan]

Two random variables & Conditional p.d.f. - III



Bayes theorem for random variables

> Combining together the two expressions for the conditional probability we get: $g(x|y) = g(y|x) \cdot \frac{f_x(x)}{f_x(y)}$

...which is nothing else that the **re-expression of the Bayes's theorem in the case of continuous r.v.s**!

 $h(y|x) \cdot f_x(x) = f(x, y) \qquad g(x|y) \cdot f_y(y) = f(x, y)$ Rewriting the same two expressions we also get: $\boldsymbol{\Sigma}$

Now we can use the definition of marginal p.d.f.s to find new expressions for them:

$$f_{x}(x) = \int_{-\infty}^{+\infty} f(x,y) dy = \int_{-\infty}^{+\infty} g(x|y) \cdot f_{y}(y) dy \qquad f_{y}(y) = \int_{-\infty}^{+\infty} f(x,y) dy = \int_{-\infty}^{+\infty} h(y|x) \cdot f_{x}(x) dx$$

...which are nothing else that the **re-expression of the Law of total probability** (slide 14 part 1A)!

Independency of events expressed as factorization for joint p.d.f.

> We have discussed earlier that: $P(A) = f_x(x)dx$ (and, in the same way, $P(B) = f_y(y)dy$). Thus, the product of the two probabilities can be expressed as:

 $P(A) \cdot P(B) = f_x(x) \, dx \cdot f_y(y) \, dy \equiv f_x(x) \, f_y(y) \, dx \, dy \quad (a)$

Let us remember now that ... two events **A** and **B** are *independent* if $P(A \cap B) = P(A) \cdot P(B)$ [*]!

From the joint pd.f. definition $P(A \cap B) = f(x, y)dx dy$ we then derive from [*] that $P(A) \cdot P(B) = f(x, y)dx dy$ (b)

Expressions (a) & (b) hold if and only if f(x, y) can be *factorized* into the product of the 2 marginal p.d.f.s:

 $f(x, y) = f_x(x) \cdot f_y(y)$

From this result: **x** and **y** can be defined as independent variables if their joint p.d.f. can be written as the product of a p.d.f. of the variable **x** times a p.d.f. of the variable **y** (specifically these p.d.f.s are the 2 marginal ones)

> Additional expressions when r.v.s are independent: $h(y|x) = \frac{f(x,y)}{f_x(x)} = \frac{f_x(x) \cdot f_y(y)}{f_x(x)} \equiv \frac{f_y(y)}{f_y(y)}$. Similarly: $g(x|y) = f_x(x)$

This means something obvious: the conditional pd.f. reduces simply to the marginal p.d.f. when the r.v.s. are independent.

CORRELATION between R.V.s

Covariance for a couple of r.v.s - I

> Let's consider 2 continuous r.v.s : (x, y). The joint p.d.f. is written as f(x, y). We can write down the following quantities:

$$\mu_x \equiv E[x] = \int_{-\infty}^{+\infty} x f(x, y) dx dy \qquad \sigma_x^2 \equiv V[x] = E[(x - \mu_x)^2]$$
$$\mu_y \equiv E[y] = \int_{-\infty}^{+\infty} y f(x, y) dx dy \qquad \sigma_y^2 \equiv V[y] = E[(y - \mu_y)^2]$$

To take into account the possible correlations among the r.v.s, that generally are not negligible and cannot be overlooked, We need to introduce a further quantity called *covariance*, defined as follows:

$$V_{xy} \equiv cov(x, y) = E[(x - \mu_x)(y - \mu_y)] = \iint_{-\infty}^{+\infty} xy \cdot f(x, y) dxdy$$

$$= E[xy - x\mu_y - y\mu_x + \mu_x\mu_y] =$$

$$= E[xy] - \mu_y E[x] - \mu_x E[y] + \mu_x\mu_y =$$

$$= E[xy] - \mu_y \mu_x - \mu_x \mu_y + \mu_x \mu_y =$$

$$= E[xy] - \mu_y \mu_x - \mu_x \mu_y + \mu_x \mu_y =$$

$$= E[xy] - \mu_y \mu_x - \mu_y \mu_x + \mu_y \mu_y =$$

$$= E[xy] - \mu_y \mu_x - \mu_y \mu_y + \mu_y \mu_y =$$

$$= E[xy] - \mu_y \mu_x - \mu_y \mu_y + \mu_y \mu_y =$$

$$= E[xy] - \mu_y \mu_x - \mu_y \mu_y + \mu_y \mu_y =$$

$$= E[xy] - \mu_y \mu_x - \mu_y \mu_y + \mu_y \mu_y =$$

$$= E[xy] - \mu_y \mu_x - \mu_y \mu_y + \mu_y \mu_y =$$

$$= E[xy] - \mu_y \mu_x - \mu_y \mu_y + \mu_y \mu_y =$$

$$= E[xy] - \mu_y \mu_x - \mu_y \mu_y + \mu_y \mu_y =$$

$$= E[xy] - \mu_y \mu_x - \mu_y \mu_y + \mu_y \mu_y =$$

Note: as expected, V_{xy} gives simply the variance V_{xx} whether the r.v.s of the pair are identical (i. e. y = x)

Covariance for a couple of r.v.s - II

We have seen (slide 15) that E[u(x, y)] can be expressed - in general - as: $E[u(x, y)] = \iint_{-\infty}^{+\infty} u(x, y) \cdot f(x, y) dx dy$... and considering the specific case of $u(x, y) = x \cdot y$: $E[xy] = \iint_{-\infty}^{+\infty} xy \cdot f(x, y) dx dy$ Wrapping up: $V_{xy} \equiv cov(x, y) = E[(x - \mu_x)(y - \mu_y)] = E[xy] - \mu_x \mu_y = \iint_{-\infty}^{+\infty} xy \cdot f(x, y) dx dy - \mu_x \mu_y$ ($\equiv \sigma_{xy}$)

Remember (see slide 16) that ... in general $E[u] \neq u(\mu)$ and thus $E[xy] \neq \mu_x \mu_y$

In conclusion: $V_{xy} = E[xy] - \mu_x \mu_y \neq 0$ (i.e. one r.v. influences the other r.v. and viceversa) Note: $V_{xy} = V_{yx}$

Since we can re-write: $\sigma_x^2 \equiv V[x] = E[(x - \mu_x)(x - \mu_x)] \equiv V_{xx}$, $\sigma_y^2 \equiv V[y] = E[(y - \mu_y)(y - \mu_y)] \equiv V_{yy}$

... it is possible to accommodate the 2 variances and the 2 (equal) covariances in a 2×2 symmetric matrix:

Covariance Matrix:

$$(V)_{xy} = \begin{pmatrix} V_{xx} & V_{xy} \\ V_{yx} & V_{yy} \end{pmatrix} = \begin{pmatrix} \sigma_x^2 & cov(x,y) \\ cov(y,x) & \sigma_y^2 \end{pmatrix} = \begin{pmatrix} E[(x-\mu_x)^2] & E[xy] - \mu_x \mu_y \\ E[xy] - \mu_x \mu_y & E[(y-\mu_y)^2] \end{pmatrix}$$

$$\equiv \begin{pmatrix} \sigma_x^2 & \sigma_{xy} \\ \sigma_{yx} & \sigma_y^2 \end{pmatrix}$$

Correlation Coefficient

With the aim to have an adimentional measure of the "degree of correlation" between the two r.v.s and it is useful to introduce the **correlation coefficient** : $cov(x, y) = V_{ry}$

$$\boldsymbol{\rho}(\boldsymbol{x},\boldsymbol{y}) = \frac{cov(x,y)}{\sigma_x \cdot \sigma_y} \equiv \frac{V_{xy}}{\sqrt{V_{xx} \cdot V_{yy}}}$$

It can be demonstrated that: $\rho(x, y) \in [-1, +1]$. We get: Maximum correlation: $\rho(x, y) = +1$ NO correlation: $\rho(x, y) = 0$ maximum anti-correlation: $\rho(x, y) = -1$

It is easy to discuss the correlation coefficient by means of these scatter plots of the r.v.s x and y:



A.Pompili (E+)

Statistical M&T for Data Analysis.

Independence & uncorrelation - I

In the every-day language - often - the physicists talk about *uncorrelated* variables implicitely implying *independent* ones, although this is not correct. We will argue - instead - that strictly speaking ...
the condition of uncorrelation is <u>weaker</u> than the condition of independency !

Indeed we will show that ... independency implies uncorrelation but the viceversa is not true!

Independence & uncorrelation - I

In the every-day language - often - the physicists talk about *uncorrelated* variables implicitely implying *independent* ones, although this is not correct. We will argue - instead - that strictly speaking ...
the condition of uncorrelation is <u>weaker</u> than the condition of independency !

Indeed we will show that ... independency implies uncorrelation but the viceversa is not true!

> To argue this, let me start recalling (see slide 27) that ...

 $\begin{bmatrix} \dots \text{ if } (x, y) \text{ are (mutually) independent random variables their joint p.d.f. factorizes: } \mathbf{f}(x, y) = \mathbf{f}_x(x) \cdot \mathbf{f}_y(y) \\ \text{and in this case: } \mathbf{E}[xy] = \iint_{-\infty}^{+\infty} xy \cdot f(x, y) dx dy = \int_{-\infty}^{+\infty} x \cdot f_x(x) dx \cdot \int_{-\infty}^{+\infty} y \cdot f_y(y) dy = \mathbf{E}[x] \cdot \mathbf{E}[y] \\ \text{which implies that : } \mathbf{V}_{xy} = \mathbf{E}[xy] - \mu_x \mu_y = \mathbf{E}[x] \cdot \mathbf{E}[y] - \mu_x \mu_y = \mu_x \mu_y - \mu_x \mu_y = \mathbf{0} \text{ (thus } \mathbf{\rho}_{xy} = \mathbf{0}) \end{aligned}$

 \Rightarrow) We have proved that : **INDEPENDENCY** \Rightarrow **UNCORRELATION**

Independence & uncorrelation - I

In the every-day language - often - the physicists talk about *uncorrelated* variables implicitely implying *independent* ones, although this is not correct. We will argue - instead - that strictly speaking ...
the condition of uncorrelation is <u>weaker</u> than the condition of independency !

Indeed we will show that ... independency implies uncorrelation but the viceversa is not true!

> To argue this, let me start recalling (see slide 29) that ...

 $\begin{bmatrix} \dots \text{ if } (x, y) \text{ are (mutually) independent random variables their joint p.d.f. factorizes: } \mathbf{f}(x, y) = \mathbf{f}_x(x) \cdot \mathbf{f}_y(y) \\ \text{and in this case: } \mathbf{E}[xy] = \iint_{-\infty}^{+\infty} xy \cdot \mathbf{f}(x, y) dx dy = \int_{-\infty}^{+\infty} x \cdot \mathbf{f}_x(x) dx \cdot \int_{-\infty}^{+\infty} y \cdot \mathbf{f}_y(y) dy = \mathbf{E}[x] \cdot \mathbf{E}[y] \\ \end{bmatrix}$ which implies that : $V_{xy} = E[xy] - \mu_x \mu_y = E[x] \cdot E[y] - \mu_x \mu_y = \mu_x \mu_y - \mu_x \mu_y = 0$ (thus $\rho_{xy} = 0$) INDEPENDENCY \Rightarrow UNCORRELATION ⇒) We have proved that : ⇐) To prove - instead - that the viceversa does not hold, i.e. **INDEPENDENCY** ⇐ **UNCORRELATION** ... we need to find at least one example characterized by dependency in spite of existing uncorrelation (next slide) $(\mathbf{v} = \mathbf{f}(\mathbf{x}))$ $(V_{rv} = 0)$

>> A suitably easy example is $(x, y) = (x, x^2)$ namely when $y = u(x) = x^2$!

To make easier the demonstration let's suppose that ... *x* is distributed symmetrically around 0, with a p.d.f. f(x), i.e.: $\mu_x \equiv E[x] = \int_{-\infty}^{+\infty} x \cdot f_x(x) dx = 0$ From the definition of variance: $\sigma_x^2 \equiv V[x] = \int_{-\infty}^{+\infty} (x - 0)^2 \cdot f_x(x) dx \equiv \int_{-\infty}^{+\infty} x^2 \cdot f_x(x) dx$

>> A suitably easy example is $(x, y) = (x, x^2)$ namely when $y = u(x) = x^2$!

To make easier the demonstration let's suppose that ...

x is distributed symmetrically around 0, with a p.d.f. f(x), i.e.: $\mu_x \equiv E[x] = \int_{-\infty}^{+\infty} x \cdot f_x(x) dx = 0$ From the definition of variance: $\sigma_x^2 \equiv V[x] = \int_{-\infty}^{+\infty} (x - 0)^2 \cdot f_x(x) dx \equiv \int_{-\infty}^{+\infty} x^2 \cdot f_x(x) dx$

Let's calculate the expectation value of the r.v. $y: \mu_y \equiv E[y] = E[u(x)] = \int_{-\infty}^{+\infty} u(x) \cdot f(x) dx = \int_{-\infty}^{+\infty} x^2 \cdot f_x(x) dx = \sigma_x^2$

Note (for completeness) that: f(x) is the marginal for x i.e. $f_x(x)$; analogously $g(u) = g(y) = g_y(y)$ would be the marginal for y.

Finally let's calculate the covariance:
$$V_{xy} = E\left[(x - \mu_x)(y - \mu_y)\right] = E\left[(x)(x^2 - \sigma_x^2)\right] = E\left[x^3 - x\sigma_x^2\right] = E\left[x^3\right] - \sigma_x^2 E\left[x\right] = E\left[x^3\right] - \sigma_x^2 E\left[x\right] = 0$$
(central moment of order-3 is null for a symmetric $f(x)$!)







Fig. 1.10 Scatter plot of random variables x and y which are not independent (i.e. $f(x, y) \neq f_x(x)f_y(y)$) but for which $V_{xy} = 0$ because of the particular symmetry of the distribution.

borrowed by G.Cowan

Note: see in-depth slides for another example.

Removing and introducing correlations by means of change variable - I

 $\boldsymbol{\Sigma}$ It is possible to remove (or introduce) a correlation by operating a change of variables, namely $(x, y) \rightarrow (x', y')$





The rotation-in-the-plane matrix:

 $A = \begin{pmatrix} \cos\vartheta & \sin\vartheta \\ -\sin\vartheta & \cos\vartheta \end{pmatrix}$

It can be calculated (G.Cowan, 1.7) that the angle is:

$$tan(2\vartheta) = \left(\frac{2V_{xy}}{\sigma_y^2 - \sigma_x^2}\right) \equiv \left(\frac{2\rho_{xy}\sigma_x\sigma_y}{\sigma_y^2 - \sigma_x^2}\right)$$

Note that the matrix A is such that the matrix $U = A \downarrow V \downarrow A^T$ is diagonal !

(I will comment further ... a few slides later)

row by column products

A.Pompili (E+)

Statistical M&T for Data Analysis.

Removing and introducing correlations by means of change variable - II

> An example of possible introduction of some correlation between two variables is a rotation in their plane as well:



Covariance for more than 2 r.v.s

> Let's consider N r.v.s: $(x_1, \dots, x_i, \dots, x_j, \dots, x_N)$

The variance of the single r.v. - regardless the others - is simply defined as:

To take into account the mutual correlations we have to introduce a coviariance for each pair (i, j):

The *N* variances and the N(N - 1) covariances (each two of them are equal by symmetry, i.e. $\sigma_{ij} = \sigma_{ji}$) can be accomodated in the **covariance matrix**, an $N \times N$ symmetric, sometimes called **error matrix**:

Note: If the covariance matrix is not positive defined... ... there must be at least one linear relationship among the r.v.s.

> A *global* correlation coefficient can be introduced when N > 2 (see in-depth slides)

 $\sigma_i^2 = E[(x_i - E[x_i])^2]$

$$V_{ij} \equiv \sigma_{ij} = E[(x_i - E[x_i])(x_j - E[x_j])]$$



Diagonalization of the covariance matrix - I

> It can be demonstrated that

... it is always possible, in the framework of linear algebra, to find an *orthogonal transformation* of $N \ge 2$ variables

 $(x_1, \dots, x_N) \rightarrow (y_1, \dots, y_N)$ for which the "new" covariance matrix for \vec{y} is diagonal while the "old" one for \vec{x} was not

It's common to say that this transformation "diagonalizes the covariance matrix", i. e. this transformation is able to remove any existing correlation.

Let's discuss this result:

- original variables & covariance matrix: (x_1, \dots, x_N) , $V_{ij} = cov(x_i, x_j)$
- transformed variables & new diagonal covariance matrix: $(y_1, ..., y_N)$, $U_{ij} = cov(y_i, y_j)$

It can be demonstrated that it is always possible to find a linear transformation, namely by means of a matrix so that each y_i is a linear combination of the (x_1, \dots, x_N) :

 $y_i = \sum_{j=1}^{N} A_{ij} x_j \quad (\forall i) \quad (\#)$ In this case the transformation matrix A is such that the new matrix $U = AVA^T$ is diagonal, and has the property that the transpose matrix coincides with the inverse ($A^T = A^{-1}$) and thus $U = AVA^{-1}$. This transformation is called *orthogonal* and it corresponds - in linear algebra - to the rotation of the vector \vec{x} into the vector \vec{y} so that the vector norm is kept constant. (see also next side)

Diagonalization of the covariance matrix - II

We can formalize what just said using the vectorial notation and the matrix formalism:

$$\vec{x} = \begin{pmatrix} x_1 \\ \cdots \\ x_N \end{pmatrix}, \quad \vec{x}^T = (x_1 \cdots x_N), \quad \vec{y} = A\vec{x} \Leftrightarrow \begin{pmatrix} y_1 \\ \cdots \\ y_N \end{pmatrix} = \begin{pmatrix} A_{11} & \cdots & A_{1N} \\ \cdots & \cdots & \cdots \\ A_{N1} & \cdots & A_{NN} \end{pmatrix} \cdot \begin{pmatrix} x_1 \\ \cdots \\ x_N \end{pmatrix}$$
$$|\vec{y}|^2 = \vec{y}^T \cdot \vec{y} = \vec{x}^T \mathbf{A}^T \cdot A \, \vec{x} = \vec{x}^T \mathbf{A}^{-1} \cdot A \, \vec{x} = \vec{x}^T \mathbf{I} \, \vec{x} = \vec{x}^T \vec{x} = |\vec{x}|^2 \text{ : vector norm is preserved}$$

$$U_{ij} = cov(y_{i}, y_{j}) \stackrel{(\#)}{=} cov\left(\sum_{k=1}^{N} A_{ik}x_{k}, \sum_{\ell=1}^{N} A_{j\ell}x_{\ell}\right) \stackrel{(\#)}{=} cov(u, v) = E[uv] - \mu_{u}\mu_{v},$$

$$E[u(x)] = u(\mu) \quad \text{IF} \quad u(x) \text{ is linear in } x$$

$$E[a_{1}u(x) + a_{2}v(x)] = a_{1}E[u(x)] + a_{2}E[v(x)]$$

$$= \sum_{k=1}^{N} \sum_{\ell=1}^{N} A_{ik} \quad A_{j\ell} \stackrel{V_{k\ell}}{=} cov(x_{k}, x_{\ell}) \stackrel{(\#)}{=} cov(x_{k}, x_{\ell}) \stackrel{(\#)}{=} cov(x_{k}, x_{\ell}) \stackrel{(\#)}{=} cov(x_{k}, x_{\ell}) \stackrel{(\#)}{=} cov(x_{k}, x_{\ell})$$

$$= \left(\sum_{k=1}^{N} \sum_{\ell=1}^{N} A_{ik} \quad V_{k\ell} \quad A^{T}_{\ell j} \stackrel{(\#)}{=} cov(x_{k}, x_{\ell}) \stackrel{(\#)}{=} cov(x_{k}, x_{\ell}) \stackrel{(\#)}{=} cov(x_{k}, x_{\ell}) \stackrel{(\#)}{=} cov(x_{k}, x_{\ell})$$

$$= \left(\sum_{k=1}^{N} \sum_{\ell=1}^{N} A_{ik} \quad V_{k\ell} \quad A^{T}_{\ell j} \stackrel{(\#)}{=} cov(x_{k}, x_{\ell}) \stackrel{(\#)}{=} cov(x_{k}, x_{\ell}) \stackrel{(\#)}{=} cov(x_{k}, x_{\ell}) \stackrel{(\#)}{=} cov(x_{k}, x_{\ell})$$

Statistical M&T for Data Analysis.

UNCERTAINTY (ERROR) PROPAGATION

Propagation of the variances - I

Suppose we have N r.v.s $(x_1, ..., x_n)$, that we can write - in a compact way - as the vector $\vec{x} \equiv (x_1, ..., x_n)$, distributed according to the joint p.d.f. $f(\vec{x})$ that we suppose is <u>not</u> fully known since we assume we only know:

- the *N* expectation values, namely the vector $\vec{\mu} \equiv (\mu_1, ..., \mu_n)$
- the $N \times N$ covariance matrix V_{ij}

Let's now consider a function $y = u(\vec{x})$ and we have seen (slides 17-18) that ...

... we can determine the p.d.f. of y - say g(u) - if we know the p.d.f $f(\vec{x})$ which, however, is not our case here! Thus, we want to determine just E[y] and V[y].

We will see that this is possible, even if we will get **approximated** (but still **useful**) expressions!

The procedure starts from the expansion in series - truncated at 1st order - of the function $y(\vec{x})$ around the vector of the expectation values $\vec{\mu}$:

$$y(\vec{x}) \cong y(\vec{\mu}) + \sum_{i=1}^{N} \left(\frac{\partial y}{\partial x_i}\right)_{\vec{x} = \vec{\mu}} \cdot (x_i - \mu_i) + \times$$

The expectation value can be easily calculated at first order:

$$\boldsymbol{E}[\boldsymbol{y}(\vec{\boldsymbol{x}})] \cong E[\boldsymbol{y}(\vec{\boldsymbol{\mu}})] + E[\sum_{i=1}^{N} \dots] \stackrel{\checkmark}{=}$$

I can apply one of the properties of the expectation value of a variable; consider that the derivatives are calculated for $\vec{x} = \vec{\mu}$ so they are just real numbers

$$= E[y(\vec{\mu})] + \sum_{i=1}^{N} \left(\frac{\partial y}{\partial x_{i}}\right)_{\vec{x}=\vec{\mu}} \cdot E(x_{i} - \vec{\mu}_{i})^{0} = E[y(\vec{\mu})]$$

(as expected: at 1st order the dependency is linear)

Propagation of the variances - II



Propagation of the variances - III

Thus, we got:
$$\sigma_{\mathbf{y}}^{\mathbf{z}} = E[y^2] - (E[y])^2 \cong \sum_{i,j=1}^{N} \left(\frac{\partial y}{\partial x_i} \frac{\partial y}{\partial x_j}\right)_{\vec{x}=\vec{\mu}} V_{ij}$$
 : equation of the error propagation
If we conventionally define the vector of partial derivatives $A = \left(\frac{\partial y}{\partial x_1}, ..., \frac{\partial y}{\partial xN}\right)$
... we can re-express this result in matrix notation: $\sigma_{\mathbf{y}}^{\mathbf{z}} = \left(\frac{\partial y}{\partial x_1}, ..., \frac{\partial y}{\partial xN}\right) \cdot \begin{pmatrix} V_{11} & ... & V_{1N} \\ ... & ... & ... \\ V_{N1} & ... & V_{NN} \end{pmatrix} \left[\begin{pmatrix} \frac{\partial y}{\partial x_1} \\ \frac{\partial y}{\partial xN} \\ \frac{\partial y}{\partial xN} \end{pmatrix} \right]$
... and more compactly: $\sigma_{\mathbf{y}}^{\mathbf{z}} = \mathbf{A} \mathbf{V} \mathbf{A}^{\mathbf{T}}$
Do not forget that ... this result is valid in the approximation in
which $y(\vec{x})$ is approximated by the Taylor expansion truncated
to the 1st order, namely in the linearity approximation around $\vec{\mu}$!
In the particular case in which the $(x_1, ..., x_n)$ are all *uncorrelated* among each other, i.e. $\begin{cases} V_{ii} = \sigma_i^2 (\forall i) \\ V_{ij} = 0 (\forall i \neq j) \\ ... then the propagation formula reduces to: $\sigma_{\mathbf{y}}^{\mathbf{z}} \equiv \sum_{i=1}^{N} \left(\frac{\partial y}{\partial xi}\right)_{\vec{x}=\vec{\mu}}^2 \sigma_i^2$ (the well-known "error propagation formula")$

A.Pompili (E+)

 $\mathbf{\Sigma}$

Statistical M&T for Data Analysis.

≠ j)

Propagation of the variances : special cases

>> Usual cases are these 4: The standard deviations sum up in guadrature **Relation between** x_1 and x_2 uncorrelated ($V_{12} = 0$) x_1 and x_2 correlated ($V_{12} \neq 0$) x_1 and x_2 $\sigma_v^2 = \sigma_{x_1}^2 + \sigma_{x_2}^2$ $\sigma_v^2 = \sigma_{x_1}^2 + \sigma_{x_2}^2 + 2V_{12}$ $y = x_1 + x_2$ $\sigma_y^2 = \sigma_{x_1}^2 + \sigma_{x_2}^2$ $\sigma_{v}^{2} = \sigma_{x_{1}}^{2} + \sigma_{x_{2}}^{2} - 2V_{12}$ $y = x_1 - x_2$ $\frac{\sigma_y^2}{(\mu_1 \cdot \mu_2)^2} = \frac{\sigma_{x_1}^2}{\mu_1^2} + \frac{\sigma_{x_2}^2}{\mu_2^2} + 2\frac{V_{12}}{(\mu_1 \cdot \mu_2)} \quad \left\{ \begin{array}{l} \mu_1 = E[x_1] \\ \mu_2 = E[x_2] \end{array} \right.$ $\frac{\sigma_y^2}{(\mu_1 \cdot \mu_2)^2} = \frac{\sigma_{x_1}^2}{\mu_1^2} + \frac{\sigma_{x_2}^2}{\mu_2^2}$ $\mathbf{y} = x_1 \cdot x_2$ $\frac{\sigma_y^2}{(\mu_1/\mu_2)^2} = \frac{\sigma_{x_1}^2}{\mu_1^2} + \frac{\sigma_{x_2}^2}{\mu_2^2} - 2\frac{V_{12}}{(\mu_1/\mu_2)} \qquad \begin{bmatrix} \mu_1 = E[x_1] \\ \mu_2 = E[x_2] \end{bmatrix}$ $\frac{\sigma_y^2}{(\mu_1/\mu_2)^2} = \frac{\sigma_{x_1}^2}{\mu_1^2} + \frac{\sigma_{x_2}^2}{\mu_2^2}$ $y = \frac{x_1}{x_2}$ (*) Very usuful because we deal very often with ratios ! (*) Another way to remember it: $\sigma_{y}^{2} = \frac{x_{1}}{r_{2}} \cong \frac{\sigma_{1}^{2}}{\mu_{2}^{2}} + \frac{\sigma_{2}^{2}}{\mu_{2}^{2}} \cdot \left(\frac{\mu_{1}^{2}}{\mu_{2}^{2}}\right) \quad ... \text{ or even better: } \sigma_{y} = y \cdot \left(\frac{\sigma_{x_{1}}^{2}}{x_{1}^{2}} + \frac{\sigma_{x_{2}}^{2}}{x_{2}^{2}}\right)$ The *relative* standard deviations sum up in quadrature (where expectation values are substituted by the actual value)

A.Pompili (E+)

Statistical M&T for Data Analysis.

PART 2B - IN-DEPTH SLIDES

A.Pompili (E+)

Statistical M&T for Data Analysis

Since all **symmetric** p.d.f.s have null odd central moments, the central moments of odd order (3, 5, ...) provide a measurement of the asymmetry of a generic distribution (remember the one of 1^{st} order is null).

In order to have an adimentional quantity we prefer divide by $\sigma_x^3 = (\mathrm{V}[x])^{3/2}$:

$$\gamma_1 = \frac{E[(x-\mu)^3]}{\sigma_x^3}$$

> For a p.d.f. characterized by a central symmetric peak, its peaking "level" (or "degree"), let's call it "**sharpness**", can be measured through :

kurtosis of a p.d.f. is defined as:

$$\gamma_2 = \frac{E[(x-\mu)^4]}{\sigma_x^4} - 3$$

This ad hoc definition derives from the aim to have $\gamma_2 = 0$ for a Gaussian p.d.f., thus this "sharpness" is compared to that of the Gaussian used as the reference.

 \triangleright Demonstrate the expression for V[x] of a mixture of sub-samples:

$$\begin{split} f(x) &= \sum_{i} \underbrace{\Phi_{i}}_{i} E_{i} \left[\left(x_{i} - \mu_{i} - \overline{\Delta_{i}} \right)^{2} \right] = \\ &= \sum_{i} \underbrace{\Phi_{i}}_{i} E_{i} \left[\left(x - \mu_{i} \right)^{2} + \overline{\delta_{i}}^{2} - 2\overline{\Delta_{i}} \left(x - \mu_{i} \right) \right] = \\ &= \sum_{i} \underbrace{\Phi_{i}}_{i} \left\{ E_{i} \left[\left(x - \mu_{i} \right)^{2} \right] + E_{i} \left[\overline{\Delta_{i}}^{2} \right] - E_{i} \left[2\overline{\Delta_{i}} \left(x - \mu_{i} \right) \right] \right\} \\ &= \sum_{i} \underbrace{\Phi_{i}}_{i} \left\{ V_{i} \left[x \right] + \overline{\delta_{i}}^{2} - 2\overline{\Delta_{i}} \left[E_{i} \left[\left(x - \mu_{i} \right) \right] \right\} \right\} \\ &= \sum_{i} \underbrace{\Phi_{i}}_{i} \left\{ V_{i} \left[x \right] + \overline{\delta_{i}}^{2} \right\} \end{split}$$

putting together I get the (overall) variance :

 $V[x] = \sum_{i} \varphi_{i} \cdot \left\{ V_{i}[x] + \left[\sum_{j \neq i} \varphi_{i} (\mu_{j} - \mu_{i}) \right]^{2} \right\}$

Now I rewrite in a useful way the deviations δ_i :

$$\begin{split} \overline{b}_{i} &= \mu - \mu_{i} = \sum_{j} \overline{\Phi}_{j} \mu_{j} + \overline{\Phi}_{i} \mu_{i} = \\ &= \sum_{j \neq i} \overline{\Phi}_{j} \mu_{j} + \overline{\Phi}_{i} \mu_{i} - \mu_{i} = \\ &= \sum_{j \neq i} \overline{\Phi}_{j} \mu_{j} - (1 - \overline{\Phi}_{i}) \mu_{i} = (\sum_{j} \overline{\Phi}_{j} = 1) \\ &= \sum_{j \neq i} \overline{\Phi}_{j} \mu_{j} - \sum_{j} \overline{\Phi}_{j} \mu_{i} + \overline{\Phi}_{i} \mu_{i} = \\ &= \sum_{j \neq i} \overline{\Phi}_{j} \mu_{j} - \sum_{j \neq i} \overline{\Phi}_{j} \mu_{i} - \overline{\Phi}_{i} \mu_{i} + \overline{\Phi}_{i} \mu_{i} = \\ &= \sum_{j \neq i} \overline{\Phi}_{j} (\mu_{j} - \mu_{i}) \end{split}$$

Additional example of dependence with uncorrelation

 $\mathbf{\Sigma}$

Example 2.7 Uncorrelated Variables May not Be Independent An example of PDF that describes uncorrelated variables that are not independent is given by the sum of four two-dimensional Gaussian PDFs as specified below:

$$f(x, y) = \frac{1}{4} \left[g(x; \mu, \sigma) g(y; 0, \sigma) + g(x; -\mu, \sigma) g(y; 0, \sigma) \right]$$

$$g(x; 0, \sigma) g(y; \mu, \sigma) + g(x; 0, \sigma) g(y; -\mu, \sigma) \right],$$
(2.83)

where g is a one-dimensional Gaussian distribution.



borrowed by L.Lista

Correlation coefficient for more than 2 r.v.s

> For each pair (i, j) a correlation coefficient can be defined in the standard way: $\rho(x_i, x_j) = \frac{v_{ij}}{\sigma_i, \sigma_i}$

Nevertheless it can be introduced a more useful indicator, the global correlation coefficient :

- take a generic the r.v. \boldsymbol{x}_{k}
- consider the correlations $\rho(x_k, y)$
- consider the linear combination y of all the other N-1 r.v.s $x_{i\neq k}$
- define the global correlation coefficient $\rho_k = max\{\rho(x_k, y)\}$

as the quantity that measures the total amount of correlation among x_k and all the others $x_{i\neq k}$

 $\rho_k = 0$ $\langle x_k \rangle$ is fully uncorrelated with all the others $x_{i \neq k}$ Thus :

 $\rho_k = 1$ $\langle m \rangle$ x_k is fully correlated with at least one linear combination of the others $x_{i \neq k}$

An useful result (given without demonstration) is the following

ng:
$$\rho_k = \sqrt{1 - [V_{kk} \cdot (V^{-1})_{kk}]^{-1}}$$

... where ... $\begin{cases} (V)_{kk} : \text{diagonal element of the covariance matrix} \\ (V^{-1})_{kk} : \text{diagonal element of the inverse of the covariance matrix} \end{cases}$

A.Pompili (E+)